

ORIGINAL PAPERS

Adv Clin Exp Med 2012, 21, 2, 207–213
ISSN 1899–5276

© Copyright by Wroclaw Medical University

RAHIME BEDIR FINDIK¹, AYSE NURCAN UNLUER², ERHAN ŞAHİN³, ÖMER FARUK BOZKURT³,
JALE KARAKAYA⁴, ALİ ÜNSAL³

Urinary Incontinence in Women and Its Relation with Pregnancy, Mode of Delivery, Connective Tissue Disease and Other Factors

Nietrzymanie moczu u kobiet i jego związek z ciążą, rodzajem porodu, chorobą tkanki łącznej i innymi czynnikami

¹ Kecioren Education and Research Hospital, Department of Obstetrics and Gynecology, Ankara, Turkey

² Bilecik Central Hospital, Department of Obstetrics and Gynecology, Bilecik, Turkey

³ Kecioren Education and Research Hospital, Department of Urology, Ankara, Turkey

⁴ Hacettepe University, Department of Biostatistics, Ankara, Turkey

Abstract

Objectives. The aim of this study was to determine the relationship between urinary incontinence (UI) and parameters such as pregnancy, mode of delivery and other factors.

Material and Methods. The study was based on a questionnaire administered to 761 patients. After their age, menopausal status, number of pregnancies, number of deliveries and history of connective tissue disease (CTD) were recorded, the data were analyzed using a χ^2 test.

Results. In patients who are post-menopausal and over 40, both stress incontinence (SI) and urge incontinence (URI) were found to be significantly higher. It was also established that both SI and URI increased with the number of pregnancies. As the number of normal vaginal deliveries increased, the rate of both SI and URI increased, while increases in the number of cesarean sections affected neither SI nor URI. Among women who had had abortions, the rate of SI was higher, and it increased as the number of abortions increase. Neither the occurrence of abortions nor increased numbers of abortions influenced the rate of URI. In patients with CTD, both SI and URI increase.

Conclusions. Urinary incontinence in women seems to be influenced by the mode of delivery, menopause, age and CTD (Adv Clin Exp Med 2012, 21, 2, 207–213).

Key words: urge incontinence, stress incontinence, urinary incontinence.

Streszczenie

Cel pracy. Określenie związków między nietrzymaniem moczu (n.m.) a wskaźnikami, takimi jak ciąża, rodzaj porodu i inne.

Materiał i metody. Badanie zostało oparte na kwestionariuszu, który wypełniło 761 pacjentek. Po odnotowaniu ich wieku, stanu menopauzalnego, liczby ciąży, liczby porodów i chorób tkanki łącznej (CTD) w wywiadzie, dane analizowano za pomocą testu χ^2 .

Wyniki. U pacjentek, które są po menopauzie i powyżej 40 lat zarówno wysiłkowe nietrzymanie moczu (w.n.m.), jak i nietrzymanie moczu spowodowane naglącymi parciami (n.m.) okazały się znacznie częstsze. Ustalono również, że częstotliwość w.n.m. i n.m. wzrasta wraz z liczbą ciąży. Wraz ze wzrostem liczby porodów siłami natury zwiększa się częstotliwość w.n.m. i n.m., podczas gdy wzrost liczby cięć cesarskich nie wpływa na częstotliwość w.n.m. i n.m. Wśród kobiet, które dokonały aborcji częstotliwość n.m. była większa; tym większa, im większa była liczba aborcji. Ani występowanie aborcji, ani ich większa liczba nie miały wpływu na częstotliwość n.m. U chorych na CTD częstotliwość w.n.m. i n.m. były większe.

Wnioski. Na występowanie nietrzymania moczu u kobiet wpływa rodzaj porodu, menopauza, wiek i CTD (Adv Clin Exp Med 2012, 21, 2, 207–213).

Słowa kluczowe: nietrzymanie moczu spowodowane naglącymi parciami, wysiłkowe nietrzymanie moczu, nietrzymanie moczu.

Urinary incontinence (UI) is defined by the International Continence Society as involuntary passing of urine [1, 2]. UI is one of the most commonly occurring conditions among women [3–6, 7]. It is more common in women than in men in all age groups [5]. Hunskar et al. [2, 8] found the prevalence of UI to be 20–30% among young adult women and 30–40% among middle-aged women [5]. It is a serious public health problem among women, and has physical, social, psychological and economic implications [9]. Most women do not report this complaint to physicians, as they feel ashamed [2, 8–9]. Although it is not a life threatening condition, UI is important since it impairs the quality of life.

The fact that UI is more common in women than in men indicates the contribution of factors such as fecundity and pregnancy to UI. The present study investigates the relation between UI and parameters such as the number of pregnancies and abortions, mode of delivery, age, menopausal status and history of connective tissue disease.

Material and Methods

A total of 761 patients from various age groups who had been referred to the Kecioren Education and Research Hospital gynecology clinic between January and September 2010 were included in this study. Patients with communication problems, mental retardation and neurological disturbances, as well as those who had previously undergone surgery for urinary incontinence were excluded from the study. Approval for the study was obtained from the Ethics Committee of Kecioren Education and Research Hospital, and all patients consented to participate in the study in accordance with the principles of the Helsinki Declaration.

For the study, each patient's name, age and menopausal/menstruation status was recorded. If the patient was post-menopausal, its duration was recorded. The patients were asked about the presence of rheumatoid arthritis (RA), lupus and other collagen and rheumatic diseases. The presence of stress incontinence (SI) was inquired into by asking if the patient passed urine when laughing, coughing or sneezing; if so, the patient was listed as SI positive. The question "do you experience an urgent need to urinate and leak then urine before you reach a toilet?" was asked to determine the presence of urge incontinence (URI). The overall number of pregnancies, overall number of deliveries, mode of delivery and the number of induced and spontaneous abortions were recorded.

The data obtained was evaluated using SPSS

software. For the descriptive statistics, percentage distribution figures were used, and in the analysis of relations, a χ^2 analysis was used.

Results

Among the total of 761 patients, SI was detected in 272 (35.7%), and URI in 312 (41%). There were 135 patients with CTD (connective tissue disease) (17.7%).

Regarding the number of pregnancies, 7.6% had never been pregnant (n = 58); 11.4% had been pregnant once (n = 87); 22.5% had had two pregnancies (n = 171); 21.9% had had three pregnancies (n = 167); 14.5% had had four pregnancies (n = 110); and 22.1% (n = 168) had had five or more pregnancies (Table 1).

See Table 1 for the frequency of the parameters investigated: urinary incontinence, age, number of pregnancies, mode of delivery, number of abortions, menopausal status and age, and Table 2 for the analysis of relations among the parameters.

It was established that higher the age, the higher the rate of SI and URI. While the rate of URI was 20.2% between the ages of 20 and 29, it was 33.5% between the ages of 30 and 39 and 52% in the group aged 40 or more. This increase was statistically significant ($p = 0$). As to the rate of SI, it was 17% in the 20–29 age group, 33.5% in the 30–39 age group and 43.8% in the 40+ age group, which represents a statistically significant difference ($p < 0.01$).

In the comparison of pre- and post-menopausal patient groups, the rate of SI was significantly higher in post-menopausal women (46.7%) than in pre-menopausal women (31.1%; $p = 0$). The rate of URI was also significantly higher in post-menopausal women (67.4%) than in pre-menopausal women (29.8%; $p = 0$).

The rate of SI and URI increased significantly with the number of pregnancies as well ($p = 0$ in both cases).

In the analysis of the relation between the mode of delivery and incontinence, a significant positive correlation was found between SI and normal vaginal delivery (NVD). As the number of normal deliveries increased, so did the rate of SI: after one NVD the rate of SI was 7.7%; after two NVDs it was 25%; after three NVDs it was 31.3%; and after four NVDs: it was 15.8% ($p = 0$). No significant relation was found between SI and the number of caesarean sections (C/S): After one C/S the SI rate was 9.9%; after two C/S it was 4.8%; after three C/S it was 0.4% ($p = 0.132$).

The analysis established that the rate of URI increased with the number of normal vaginal

Table 1. The frequency of the characteristics analysed**Tabela 1.** Częstotliwość występowania badanych cech

Characteristic (Cecha)	Category (Kategoria)	Percent and number (N) (Odsetek i liczba)
Urinary incontinence (UI) (Nietrzymanie moczu)	stress incontinence (SI) urge incontinence (URI)	35.7% (N:272) 41% (N:312)
CTD (Connective tissue disease) (Choroba tkanki łącznej)	yes no	17.7% (N:135) 82.3% (N:626)
Number of pregnancies (G) (Liczba ciąży)	0G 1G 2G 3G 4G 5G	7.6% (N:58) 11.4% (N:87) 22.5% (N:171) 21.9% (N:167) 14.5% (N:110) 22.1% (N:168)
The number of cesarean sections (C/S) (Liczba cięć cesarskich)	1C/S 2C/S 3C/S 4C/S	9.7% (N:74) 6.8% (N:52) 1.4% (N:11) 0.1% (N:1)
The number of vaginal deliveries (NVD) (Liczba porodów siłami natury)	1NVD 2NVD 3NVD 4NVD	12.4% (N:94) 28.6% (N:218) 22.6% (N:172) 9.5% (N:72)
The number of abortions (A) (Liczba aborcji)	1A 2A 3A 4A	17% (N:129) 10.5% (N:80) 3.9% (N:30) 1.2% (N:9)
Age (Wiek)	20–29 30–39 40 and above	21.5% (N:163) 23.2% (N:176) 52% (N:418)
Menopausal status (Stan menopauzalny)	pre-menopausal post-menopausal	70.2% (N:531) 29.8% (N:227)

deliveries up to three NVD: After one NVD the URI rate was 8.7%; after two NVDs it was 27.2%; after three NVDs it was 27.6%. After four NVDs the rate dropped to 13.1%; after five NVDs it was 6.1%. No significant relation was found between URI and cesarean delivery: After one C/S the URI rate was 8%; after two C/S it was 4.5%; after three C/S it was 1% ($p = 0.066$).

When the rate of SI was compared between the C/S-only group and the NVD+C/S groups, the rates of SI were found to be 5.5% and 9% respectively, which represents a statistically significant difference ($p = 0$). The URI rate was 6.45% in the C/S-only group and 7.15% in the NVD+C/S group, which is also a statistically significant difference ($p < 0.01$).

The rate of SI was 30.6% in the group that had never had an abortion and 45.7% in the group that had had at least one abortion, which represents a statistically significant difference ($p = 0$). As the number of abortions increased, the rate of SI increased as well ($p = 0$). The URI rate was

41.7% among women who had had no abortions and 39.5% among those who had had at least one abortion. This difference was not statistically significant ($p = 0.557$). The rate of URI did not increase with the number of abortions.

In the analysis of the relation between CTD and urinary incontinence, the rate of SI was found to be 33.9% in women without CTD and 44.4% in those with CTD, which is a statistically significant difference ($p = 0.020$). The rate of URI was 39.6% in the group without CTD and 48.9% in the group with CTD, which is also a statistically significant difference ($p = 0.040$).

Discussion

Urinary incontinence is a serious public health problem which is quite common in women. It markedly impairs quality of life and leads to physical, social and psychological problems even though is not life threatening [7, 9].

Table 2. Results of the statistical analysis**Tabela 2.** Wyniki analizy statystycznej

Characteristic (Cecha)	Category (Kategoria)	SI (%)	P	URI (%)	P
Age – years (Wiek – lata)	20–29 30–39 40 and above	17 33.5 43.8	0	20 33.5 52	0
Menopausal status (Stan menopazualny)	pre-menopausal post-menopausal	31.1 46.7	0	29.8 67.4	0
The number of pregnancies (G) (Liczba ciąży)	0G 1G 2G 3G 4G 5G or more	15.5 19.5 19.9 39.5 51.8 54	0	24.1 33.3 31.6 44.9 48.2 45.2	0
The number of vaginal deliveries (NVD) (Liczba porodów siłami natury)	1NVD 2NVD 3NVD 4NVD 5NVD	7.7 25 31.5 15.8	0	8.7 27.2 27.6 13.1 6.1	0
The number of cesarean sections (C/S) (Liczba cięć cesarskich)	1C/S 2C/S 3C/S	9.9 4.8 0.4	0.132	8 4.5 1	0.066
C/S+NVD (Cięcie cesarskie + siłami natury) C/S only (Tylko cięcie cesarskie)	yes yes	9.6 5.5	0	7.1 6.4	0
Abortions (A) (Liczba aborcji)	no yes	30.6 45.7	0	41.7 39.5	0.712
Connective tissue disease (Choroba tkanki łącznej)	yes no	64.4 33.9	0.02	48.9 39.6	0.040

Urinary incontinence is more common in women than in men [2]. Based on the idea that factors associated with fecundity (pregnancy, delivery, abortion, menopause) may be associated with this, the authors investigated the relation between the aforementioned factors, as well as connective tissue disease, and urinary incontinence in a total of 761 patients.

It is well known that incontinence increases with advancing age and menopause [10, 11]. The weakening of the muscles in the pelvic floor (as in the rest of the body) with increasing age may cause this. In this respect, the results of the current study are congruent with those reported in the literature: The investigation showed that both SI and URI rates in women increase with advancing age ($p = 0$). In addition, in menopausal women, the rate of URI was significantly higher than that in pre-menopausal women. Moreover, when the rates of SI and URI were compared in menopausal patients, URI was established to be more common than SI, which indicates that urge incontinence may become more common with menopause.

The current study also established that the rates of both SI and URI increased with increas-

ing number of pregnancies ($p = 0$ in both cases). These results are consistent with the literature and confirm that pregnancy is a risk factor irrespective of the mode of delivery [12, 13]. With each pregnancy, a woman's body is exposed to reproductive hormones to a greater degree, and some of these hormones – especially relaxin – increase urinary incontinence due to the relaxation of connective tissue [14, 15]. During pregnancy, the rate of urinary incontinence increases, independent of infection [10, 16–21]. Although a previous study suggested that the first birth has the most impact on urinary continence, in the present study it was determined that the rate of urinary continence has a positive correlation with the number of pregnancies [22, 23]. Even if they do not culminate in delivery, each increase in the number of pregnancies increases the rate of UI. In some previous studies, the relation between mode of delivery and urinary incontinence was investigated and it was determined that in some cases vaginal delivery increased the rate of urinary incontinence [12, 23–31]. In the present study, the relations between mode of delivery and URI and SI were analyzed separately. Although a signifi-

cant increase in SI was established with increasing numbers of pregnancies ($p = 0$), there was no significant relation between SI and caesarean delivery ($p = 0.132$). The same was true for URI. As the number of normal deliveries increased, the rate of URI increased as well ($p = 0$), while the number of C/S did not have any such effect ($p = 0.066$), which is consistent with the literature [23–31]. These results suggest that pelvic floor impairment may occur during normal delivery [12, 32–34]. In the Norwegian EPINCONT study, the risk of URI was found to be higher in normal vaginal delivery [24]. Urinary leaks immediately after birth are a risk factor for the development of long-term incontinence [12, 35]. In the present study, cesarean section seems to offer protection from urinary incontinence; when elective C/S, post-labor C/S and pushing/expulsion C/S were compared, the risk of urinary continence was found to be the same in all of them [23]. The later stages of the fetus's descent through the vagina during delivery seem to cause the major negative effect on continence, possibly because of connective tissue damage due to episiotomy [23]. Hypoxic muscle damage due to mechanical pressure may also play a role [23]. When muscle-relaxing factors, which are already increased during pregnancy, are added to all of these, incontinence following vaginal delivery seems inevitable. All of these factors should be investigated in more detail.

In the present study, there was a group of women who had delivered both vaginally and by C/S. In the C/S-only group, the rate of SI was found to be 5.5%, while in the C/S+NVD group it was 9.6% – a significant difference ($p = 0$). The same was true for URI. These results suggest that in the patient group with two modes of delivery, both types of incontinence occur at significantly higher rates. This means the longterm effects of vaginal delivery after C/S are open to debate. After undergoing C/S once, vaginal delivery does not seem to be very reasonable for the woman's comfort.

In addition, in the present study, patients who had undergone induced or spontaneous abortion were compared with those who had not. In the abortion group, the risk of SI was found to be significantly higher than among patients who had had no abortions ($p = 0$). In the rate of URI, no difference was found between the groups ($p = 0.557$). In addition, there was a correlation between the

number of abortions and the rate of SI ($p < 0$), but the rate of URI was not affected ($p = 0.712$).

The increase in the rate of SI with an increasing number of abortions may be due to the increase in the number of pregnancies, which suggests that pregnancy by itself may have an adverse effect on urinary incontinence [12]. The negative effect of induced abortion on the pelvic floor may also be influential.

In the present study, the effect of the presence of connective tissue disease on the rate of urinary incontinence was also investigated, which is an issue that has not been addressed before. SI was found to be significantly higher in women with connective tissue disease ($p = 0.020$); so was URI ($p = 0.040$). These results suggest that pelvic ligaments may be weaker in patients with connective tissue disease.

The authors concluded that urinary incontinence is quite an important clinical condition which influences women's quality of life. Its incidence increases with age, menopausal status, increasing number of pregnancies, the number of normal vaginal deliveries, the number of abortions, and also in the presence of connective tissue disease. Patients with these risk factors can be helped through exercise programs strengthening the pelvic floor. This is an important issue for women's health.

The present study emphasized some issues which had not previously been highlighted. One of them is the increase in the risk of urinary incontinence among women who have had both cesarean and normal deliveries. Although spontaneous vaginal delivery after a cesarean section seems to be useful for the patient's comfort in the short term, the risk/benefit ratio is debatable when the effect of pelvic floor damage on urinary incontinence is considered. This is especially worth attention given that an increase in the number of C/S does not increase the rate of incontinence.

In addition, patients with connective tissue disease may also usefully be included exercise programs for pelvic floor strengthening in addition to undergoing specific treatment.

The results of the present study are consistent with the literature with regard to pregnancy and delivery, but further studies with larger patient populations are required in order to clarify the effect of connective tissue disease and induced/spontaneous abortion on urinary incontinence.

References

- [1] Abrams P, Cardozo L, Fall M, Griffiths D, Rosier P, Ulmsten U, Van Kerrebroeck P, Victor A, Wein A: Standardisation Sub-Committee of the International Continence Society. The standardisation of terminology in lower urinary tract function: report from the standardisation sub-committee of the International Continence Society. *Urology* 2003 Jan, 61(1), 37–49. Review

[2] **Boeun Y-JS:** Predictive Risk Factors for Impaired Quality of Life in Middle-Aged Women with Urinary Incontinence. *Int Neurourol J* 2010, 14, 250–255.

[3] **Hunskaar S, Lose G, Sykes D, Voss S:** The prevalence of urinary incontinence in women in four European countries. *BJU Int* 2004, 93(3), 324–330.

[4] **Hannestad YS, Rortveit G, Sandvik H, Hunskaar S:** A community-based epidemiological survey of female urinary incontinence: the Norwegian EPINCONT study. *Epidemiology of Incontinence in the County of Nord-Trøndelag. Norwegian EPINCONT study. Epidemiology of Incontinence in the County of Nord-Trøndelag.* *J Clin Epidemiol* 2000, 53(11), 1150–1157.

[5] **Hampel C, Wienhold D, Benken N, Eggersmann C, Thüroff JW:** Prevalence and natural history of female incontinence. *Eur Urol* 1997, 32 Suppl 2, 3–12.

[6] **Chiarelli P, Brown WJ:** Leaking urine in Australian women: prevalence and associated conditions. *Women Health* 1999, 29(1), 1–13.

[7] **Wesnes SL, Hunskaar S, Bo K, Rortveit G:** The effect of urinary incontinence status during pregnancy and delivery mode on incontinence postpartum. A cohort study. *BJOG* 2009, 116(5), 700–707. Epub 2009 Feb 10.

[8] **Hunskaar S, Burgio K, Clark A, Lapitan MC, Nelson R, Sillen U et al.:** Epidemiology of faecal and urinary incontinence and pelvic organ prolapse. In: *Incontinence*. Eds.: Abrams P, Cardozo L, Khoury S, Wein A. Plymouth: Health Publication Ltd, 2005, p. 257–312.

[9] **Kocak I, Okyay P, Dundar M, Erol H, Beser E:** Female urinary incontinence in the west of Turkey: prevalence, risk factors and impact on quality of life. *Eur Urol* 2005, 48(4), 634–641.

[10] **Sharma JB, Aggarwal S, Singhal S, Kumar S, Roy KK:** Prevalence of urinary incontinence and other urological problems during pregnancy: a questionnaire based study. *Arch Gynecol Obstet* 2009, 279(6), 845–851. Epub 2008 Nov 19.

[11] **Baydock SA, Flood C, Schulz JA, MacDonald D, Esau D, Jones S, Hiltz CB:** Prevalence and risk factors for urinary and fecal incontinence four months after vaginal delivery. *J Obstet Gynaecol Can* 2009, 31(1), 36–41.

[12] **Huebner M, Antolic A, Tunn R:** The impact of pregnancy and vaginal delivery on urinary incontinence. *Int J Gynaecol Obstet* 2010, 110(3), 249–251. Epub 2010 Jun 8.

[13] **Foldspang A, Hvidman L, Mommsen S, Nielsen JB:** Risk of postpartum urinary incontinence associated with pregnancy and mode of delivery. *Acta Obstet Gynecol Scand* 2004, 83(10), 923–927.

[14] **Tincello DG, Teare J, Fraser WD:** Second trimester concentration of relaxin and pregnancy related incontinence. *Eur J Obstet Gynecol Reprod Biol* 2003, 106, 237–238.

[15] **Kristiansson P, Samuelsson E, von Schoultz B, Svardsudd K:** Reproductive hormones and stress urinary incontinence in pregnancy. *Acta Obstet Gynecol Scand* 2001, 80(12), 1125–1130.

[16] **Hvidman L, Hvidman L, Foldspang A, Mommsen S, Bugge Nielsen J:** Correlates of urinary incontinence in pregnancy. *Int Urogynecol J Pelvic Floor Dysfunct* 2002, 13(5), 278–283.

[17] **Wesnes SL, Rortveit G, Bø K, Hunskaar S:** Urinary incontinence during pregnancy. *Obstet Gynecol* 2007, 109(4), 922–928.

[18] **Sampselle CM, DeLancey JOL, Ashton-Miller J:** Urinary incontinence in pregnancy and postpartum. *Neurourol Urodyn* 1996, 15, 329–330.

[19] **Viktrup L, Lose G, Rolff M, Barfoed K:** The symptom of stress incontinence caused by pregnancy or delivery in primiparas. *Obstet Gynecol* 1992, 79(6), 945–949.

[20] **Hvidman L, Foldspang A, Mommsen S, Nielsen JB:** Postpartum urinary incontinence. *Acta Obstet Gynecol Scand* 2003, 82(6), 556–563.

[21] **Glazener CM, Herbison GP, MacArthur C, Lancashire R, McGee MA, Grant AM, Wilson PD:** New postnatal urinary incontinence: obstetric and other risk factors in primiparae. *BJOG* 2006, 113(2), 208–217.

[22] **MacLennan AH, Taylor AW, Wilson DH, Wilson D:** The prevalence of pelvic floor disorders and their relationship to gender, age, parity and mode of delivery. *BJOG* 2000, 107(12), 1460–1470.

[23] **Boyles SH, Li H, Mori T, Osterweil P, Guise JM:** Effect of mode of delivery on the incidence of urinary incontinence in primiparous women. *Obstet Gynecol* 2009, 113(1), 134–141.

[24] **Van Brummen HJ, Bruinse HW, van de Pol G, Heintz AP, van der Vaart CH:** The effect of vaginal and cesarean delivery on lower urinary tract symptoms: what makes the difference? *Int Urogynecol J Pelvic Floor Dysfunct* 2007, 18(2), 133–139. Epub 2006 Apr 21.

[25] **Viktrup L:** The risk of lower urinary tract symptoms five years after the first delivery. *Neurourol Urodyn* 2002, 21(1), 2–29.

[26] **Snooks SJ, Swash M, Mathers SE, Henry MM:** Effect of vaginal delivery on the pelvic floor: a 5-year follow-up. *Br J Surg* 1990, 77(12), 1358–1360.

[27] **Tincello DG, Adams EJ, Richmond DH:** Antenatal screening for postpartum urinary incontinence in nulliparous women: a pilot study. *Eur J Obstet Gynecol Reprod Biol* 2002, 101(1), 70–73.

[28] **Meyer S, Schreyer A, De Grandi P, Hohlfeld P:** The effects of birth on urinary continence mechanisms and other pelvic-floor characteristics. *Obstet Gynecol* 1998, 92(4 Pt 1), 613–618.

[29] **Dolan LM, Hosker GL, Mallett VT, Allen RE, Smith AR:** Stress incontinence and pelvic floor neurophysiology 15 years after the first delivery. *BJOG* 2003, 110(12), 1107–1114.

[30] **Cardozo L, Cutner A:** Lower urinary tract symptoms in pregnancy. *Br J Urol* 1997, 80 Suppl 1, 14–23. Review.

[31] **Rortveit G, Daltveit AK, Hannestad YS, Hunskaar S:** Vaginal delivery parameters and urinary incontinence: the Norwegian EPINCONT study. *Am J Obstet Gynecol* 2003, 189(5), 1268–1274.

- [32] Altman D, Ekström A, Forsgren C, Nordenstam J, Zetterström J: Symptoms of anal and urinary incontinence following cesarean section or spontaneous vaginal delivery. *Am J Obstet Gynecol* 2007, 197(5), 512.e1-7.
- [33] Burgio KL, Zyczynski H, Locher JL, Richter HE, Redden DT, Wright: Urinary incontinence in the 12-month postpartum period. *Obstet Gynecol* 2003, 102(6), 1291-1298.
- [34] Hannah ME, Hannah WJ, Hodnett ED, Chalmers B, Kung R, Willan A, Amankwah K, Cheng M, Helewa M, Hewson S, Saigal S, Whyte H, Gafni A: Term Breech Trial 3-Month Follow-up Collaborative Group. Outcomes at 3 months after planned cesarean vs planned vaginal delivery for breech presentation at term: the international randomized Term Breech Trial. *JAMA* 2002, 287(14), 1822-1823.
- [35] Viktrup L, Lose G, Rolff M, Barfoed K: The symptom of stress incontinence caused by pregnancy or delivery in primiparas. *Obstet Gynecol* 1992, 79(6), 945-949.

Address for correspondence:

Rahime Bedir Findik
Kecioren Education and Research Hospital
Department of Obstetrics and Gynecology
Kecioren, Ankara
Turkey
Tel.: +90 50503166532
E-mail: drbedir75@yahoo.com

Conflict of interest: None declared

Received: 24.07.2011

Revised: 12.12.2011

Accepted: 29.03.2012