

DIDACTICS OF MATHEMATICS

7(11)



The Publishing House
of the Wrocław University of Economics
Wrocław 2010

Editors
Janusz Łyko
Antoni Smoluk

Referee
Marian Matłoka
(Uniwersytet Ekonomiczny w Poznaniu)

Proof reading
Agnieszka Flasińska

Setting
Elżbieta Szlachcic

Cover design
Robert Mazurczyk

Front cover painting: W. Tank, *Sower*
(private collection)

© Copyright by the Wrocław University of Economics
Wrocław 2010

PL ISSN 1733-7941

Print run: 200 copies

TABLE OF CONTENTS

MAREK BIERNACKI <i>Applications of the integral in economics. A few simple examples for first-year students [Zastosowania całki w ekonomii].....</i>	5
PIOTR CHRZAN, EWA DZIWOK <i>Matematyka jako fundament nowoczesnych finansów. Analiza problemu na podstawie doświadczeń związanych z uruchomieniem specjalności Master Program Quantitative Asset and Risk Management (ARIMA) [Mathematics as a foundation of modern finance]</i>	15
BEATA FAŁDA, JÓZEF ZAJĄC <i>Algebraiczne aspekty procesów ekonomicznych [Algebraical aspects of economics processes].....</i>	23
HELENA GASPARS-WIELOCH <i>How to teach quantitative subjects at universities of economics in a comprehensible and pleasant way? [Jak uczyć ilościowych przedmiotów na uczelniach ekonomicznych w zrozumiały i przyjemny sposób?]</i>	33
DONATA KOPAŃSKA-BRÓDKA <i>Wspomaganie dydaktyki matematyki narzędziami informatyki [Information technology supporting mathematical education].....</i>	49
PATRYCJA KOWALCZYK, WANDA RONKA-CHMIELOWIEC <i>Metody matematyczne w dydaktyce ubezpieczeń na studiach ekonomicznych [Mathematical methods in the didactics of insurance on economic studies].....</i>	59
LUDOMIR LAUDAŃSKI <i>The art of conjecturing (Ars Conjectandi). On the historical origin of normal distribution [Rodowód rozkładu normalnego].....</i>	67
JANUSZ ŁYKO, ANDRZEJ MISZTAŁ <i>Wpływ zmiany liczby godzin zajęć na wyniki egzaminu z matematyki na kierunkach ekonomicznych [The impact of changes in the number of hours of classes on exam results in mathematics at the economic faculties].....</i>	81
KRZYSZTOF MAŁAGA <i>Matematyka na usługach mikroekonomii [Mathematics on microeconomics services]</i>	93
WOJCIECH RYBICKI <i>Kilka powodów, dla których opowiadamy studentom ekonomii o macierzach [Some reasons for which we tell students of economics about matrices]</i>	109
ANDRZEJ WILKOWSKI <i>On changing money and the birthday paradox [O rozmiennianiu pieniędzy i paradoksie urodzin]</i>	127
HENRYK ZAWADZKI <i>Mathematica® na usługach ekonomii [Mathematica® at economics service]</i>	135

**ON CHANGING MONEY
AND THE BIRTHDAY PARADOX****Andrzej Wilkowski**

Abstract. The paper deals with two problems of discrete probability theory that can be included to a curriculum in mathematics or statistics at universities of economics. The first one addresses the question that is interesting for economists: in how many ways one can change a certain amount of money given a fixed set of denominations. One can answer the question using either theoretical or software tools (such as Matlab or Mathematica). The other problem deals with the birthday paradox and its generalization. It is an example when intuition fails.

Keywords: birthday paradox, generating function, partition.

1. First, we will discuss several facts connected with partitions of natural numbers. A number of all partitions of a natural number n is denoted by p_n . Partitions that have different order of elements are considered to be equal. Hence, it is easy to discover that a sequence (p_n) with $n \in N$ increases at a fast pace because, for instance, $p_{200} = 3972999029388$, and p_{14031} has 127 decimal figures (Narkiewicz 1977).

An effective method to analyse sequences of numbers is to work with infinite power series of complex numbers that produce a given sequence, i.e., with generating functions. We consider them as formal series that may be added, multiplied, differentiated, etc. They are a good representation of their coefficients. Their convergence is not typically relevant (unless asymptotic analysis of coefficients is needed). In the case of partitions we denote a generating function by $P(z)$. We have

$$P(z) = 1 + \sum_{n=1}^{\infty} p_n z^n, \quad z \in \mathbb{C}.$$

Andrzej Wilkowski

Department of Mathematics, Wrocław University of Economics, ul. Komandorska 118/120,
53-345 Wrocław, Poland.

e-mail: andrzej.wilkowski@ue.wroc.pl

It is known (Palka, Ruciński 1998) that if $|z| < 1$, then

$$P(z) = \prod_{n=1}^{\infty} (1 - z^n)^{-1},$$

where the right-hand side product is absolutely and almost uniformly convergent. While comparing generating functions, it is easily shown that the number of partitions into distinct parts is equal to the number of partitions into odd parts. The respective generating functions are given by:

$$R(z) = (1 + z)(1 + z^2) \dots$$

The first parenthesis in the above product means that 1 appears in the partition once or not at all; the second parenthesis – that 2 appears once or never, etc. For partitions into odd parts, we have:

$$N(z) = (1 + z^2 + z^3 + \dots)(1 + z^3 + z^6 + \dots)(1 + z^5 + z^{10} + \dots) \dots$$

The interpretation of respective terms in parentheses is similar as in the case of the function $R(z)$. It is sufficient now to replace a term $1 + z^k$ in the generating function $R(z)$ by

$$\frac{1 - z^{2k}}{1 - z^k}.$$

To obtain the equality $R(z) = N(z)$.

2. We now focus on restricted partitions, i.e., a natural number n is to be written as a sum of predetermined parts. They are a useful tool to analyse monetary systems (Smoluk 1998). One of the first European monetary systems was created by an Athenian lawmaker, Solon (it was also a system of weights and measures), who derived it from the Babylonian number system based on sixty (threescore). Since about 594 BC, a Greek, or Attic talent was equal to 60 minas or 3600 shekels or 6000 drachmae (its weight was about 25.9 kilograms). After the coinage reform, one drachma coin contained 4.37 grams of silver. Another circulating coin was the obol whose name stemmed from a long thin metal rod used as a spit. Six obols were worth one drachma. Next, there was a stater (literally “weight”) minted of gold, electrum or silver. A gold and electrum stater was worth twenty drachmae, while a silver one – four drachmae (Hammond 1994). Let assume

an obol as a basic coin. Then the Greek monetary system is described by four generating functions:

$$A_i(z) = 1 + z^i + z^{2i} + z^{3i} + \dots,$$

where $i \in \{1, 6, 24, 120\}$ (Wilkowski 1998). Let $A(z) = 1 + a_1z + a_2z^2 + \dots$ denote a product of the four above power series. In order to determine in how many ways one could pay, say, 3600 obols (that was a price of a good slave around 400 BC), (Mizerski 1996), we have to obtain the coefficient a_{3600} of series $A(z)$. It is the number of all feasible partitions of the number 3600 in the Greek monetary system. We may use available numeric software packages such as Matlab or Mathematica. A derivative of the 3600th order of the function $A(z)$ at zero is to be calculated and divided by the factorial 3600!

It is well-known that good money is strong and stable. However, all currencies tend to lose their real purchasing power, e.g., between the 9th and the 20th centuries the British pound's loss of purchasing power was as 1:3360, or in other words, the British pound in 1988 was worth merely 0.03 per cent of its original value (Żabiński 1989). Perhaps, the euro will be such a currency, since the GDP of the European Union's countries in 2009 exceeded that of the US. The euro may also become, in addition to the US dollar, a global reserve currency. Likewise, Athenian drachmae, deniers of Charlemagne, Florentine florins, Dutch guilders, or finally, Reichsthaler played such a role in the past.

The euro, just like the majority of other monetary systems globally, is based on a decimal system. One euro is divided into 100 cents. We have got coins in 1, 2, 5, 10, 20 and 50 cent denominations. The coins are issued in 1 and 2 euro denominations as well (since 2005, the European Commission has been thinking about the introduction of banknotes in these denominations, but no final decision was reached so far), while banknotes are issued in €5, €10, €20, €50, €100, €200 and €500.

Problem (Wilkowski 2000). In how many ways one can change a 10 euro note using only 1, 5, 10 and 50 cent coins?

Solution I (theoretical). The generating function for this situation has the following form:

$$A(z) = (1 + z + z^2 + \dots)(1 + z^5 + z^{10} + \dots)(1 + z^{10} + z^{20} + \dots)(1 + z^{50} + z^{100} + \dots).$$

In other words:

$$A(z) = \frac{1}{1-z} \cdot \frac{1}{1-z^5} \cdot \frac{1}{1-z^{10}} \cdot \frac{1}{1-z^{50}}.$$

We have to determine the coefficient a_{1000} of the term z^{1000} of the function $A(z)$. It is seen that

$$A(z) = (1 + z + z^2 + z^3 + z^4)B(z^5),$$

where

$$B(z) = \frac{1}{1-z} \cdot \frac{1}{1-z} \cdot \frac{1}{1-z^2} \cdot \frac{1}{1-z^{10}}.$$

Since each term in the denominator of the function $B(z)$ is a divisor of $1 - z^{10}$, hence we get:

$$B(z) = \frac{W(z)}{(1-z^{10})^4}.$$

A polynomial $W(z)$ has the following form:

$$\begin{aligned} W(z) &= (1 + z + \dots + z^9)^2(1 + z^2 + \dots + z^8) = \\ &= 1 + 2z + 4z^2 + 6z^3 + 9z^4 + 12z^5 + 16z^6 + 20z^7 + 25z^8 + 30z^9 + \\ &\quad + 33z^{10} + 36z^{11} + 37z^{12} + 38z^{13} + 37z^{14} + \dots + 2z^{25} + z^{26}, \\ &\quad (w_i = w_{26-i}, \quad i = 0, 1, \dots, 26). \end{aligned}$$

It is known (Graham, Knuth, Patashnik 1998) that

$$\frac{1}{(1-z^{10})^4} = \sum_{n=0}^{\infty} \binom{n+3}{3} z^{10n},$$

therefore

$$B(z) = \left(\sum_{n=0}^{\infty} \binom{n+3}{3} z^{10n} \right) W(z).$$

We can now calculate the coefficient b_n appearing with z^n in the series $B(z)$. With $n = 10k + i$ and $0 \leq i < 10$, we have:

$$b_{10k+i} = w_i \binom{k+3}{3} + w_{i+10} \binom{k+2}{3} + w_{i+20} \binom{k+1}{3},$$

where w_i is the coefficient of the polynomial

$$W(z) = w_0 + w_1z + \dots + w_{26}z^{26}.$$

Because $a_{50k} = b_{10k}$ therefore

$$a_{1000} = \binom{23}{3} + 33 \binom{22}{3} + 16 \binom{21}{3} = 73871.$$

With 1, 5, 10 and 50 cent coins available, we can change a 10 euro banknote in 73871 ways.

Solution II (numerical). Using Matlab or Mathematica, we obtain the solution:

$$a_{1000} = \frac{A(0)^{(1000)}}{1000!} = 73871.$$

To conclude this section, let us note that in view of the ongoing development of cashless payments, increasing usage of credit, charge and debit cards and the like, perhaps the banknotes of high denominations will become redundant. However, it is less likely that all coins and notes will completely disappear along with current monetary systems that we study by means of generating functions.

3. When solving some probabilistic problems, we face the failure of intuition. Such problems include among others: the Chebyshev problem, the Bertrand paradox (Wilkowski 2007), the Penney's game or the Monty Hall paradox (Dniestrzański, Wilkowski 2008). This section deals with this type of a problem, i.e., the birthday paradox. In this problem we ask a question: what is the smallest number n of people that at least one pair of them will have the same birthday with the probability greater than 0.5? Students of many US universities were asked this question, with an average answer 385 (Nikodem 2010), thus proving that their intuition failed (it is interesting what the outcome would be among Polish students).

Let us therefore solve the problem contained in this question. We assume that birthdays are uniformly distributed on the set $\{1, \dots, 365\}$ (we disregard leap years, twins and seasonal variation of births). These factors do not significantly affect the solution. Elementary events are now k -term sequences whose elements are from the set $\{1, \dots, 365\}$. There are 365^k

such sequences. Let p_k denote the probability of the complement of the event under study, i.e., the event that among the k persons each one has a different birthday. Then, it holds:

$$p_k = \frac{365 \cdot 364 \cdot \dots \cdot (365 - k + 1)}{365^k} = \left(1 - \frac{1}{365}\right) \cdot \dots \cdot \left(1 - \frac{k-1}{365}\right),$$

for $2 \leq k \leq 365$ and

$$p_k = 0$$

for $k > 365$.

Using the inequality $1 + x \leq e^x$ that is true for any real number x , we obtain the following approximation:

$$p_k \leq e^{-\frac{1}{365}} \cdot \dots \cdot e^{-\frac{k-1}{365}} = e^{-\frac{k(k-1)}{730}}.$$

The smallest natural k with $p_k \leq \frac{1}{2}$ is equal 23. Using a software package such as Matlab or Mathematica yields the following approximation:

$$p_{23} \approx 0,492703, \quad p_{22} \approx 0,524305.$$

Hence, if there are at least 23 persons in a group, then a probability that at least two of them have the same birthday is greater than $\frac{1}{2}$.

Let us now consider a more difficult problem (assuming, as before, the uniformity, and disregarding leap years, twins and triplets). What is the smallest number of persons so that the probability that at least three of them share a birthday is greater than 0.5? The probability of the complementary event equals:

$$Pr(\text{no more than two out of } k \text{ persons share a birthday}) = \frac{f_k}{365^k},$$

where

$$f_k = \sum_{l \geq 0} \binom{365}{k-l} \binom{k-l}{l} \frac{k!}{2^l} \quad (\text{Adamaszek 2010}).$$

We seek the smallest natural number k that satisfies the inequality:

$$1 - \frac{f_k}{365^k} > \frac{1}{2}.$$

Using the software package Matlab or Mathematica, we obtain:

$$1 - \frac{f_{88}}{365^{88}} = 0,511\dots$$

We conclude that at least 88 persons are needed for the probability that at least three of them have the same birthday to exceed $\frac{1}{2}$. During class sessions, one can solve similar problems requesting the four, five and more persons sharing the same birthday.

Literature

- Adamaszek M. (2010). *Potrójne urodziny* [Triple birthday]. "Delta" No. 4. Wydawnictwo Uniwersytetu Warszawskiego. Warszawa.
- Dniestrzański P., Wilkowski A. (2008). *O paradoksie Halla i rzucaniu monetą* [On the Hall paradox and tossing a coin]. "Didactics of Mathematics" No. 5 (9). Wydawnictwo Akademii Ekonomicznej. Wrocław.
- Graham R.L., Knuth D.E., Patashnik O. (1998). *Matematyka konkretna* [Concrete mathematics]. Wydawnictwo Naukowe PWN. Warszawa.
- Hammond N.G.L. (1994). *Dzieje Grecji* [A history of Greece]. Państwowy Instytut Wydawniczy. Warszawa.
- Mizerski W. (1996). *Tablice historyczne* [Historical tables]. Wydawnictwo Adamantan. Warszawa.
- Narkiewicz W. (1977). *Teoria liczb* [Number theory]. Wydawnictwo Naukowe PWN. Warszawa.
- Nikodem T. (2010). *Paradoks dni urodzin i pokrewne, czyli o pewnych zagadnieniach związanych z rozmieszczeniem kul w komórkach* [Birthday paradox and the akin problems or some issues of how to allocate balls to cells]. "Delta" No. 4. Wydawnictwo Uniwersytetu Warszawskiego. Warszawa.
- Palka Z., Ruciński A. (1998). *Wykłady z kombinatoryki. Część I* [Lectures in combinatorics. Part I]. Wydawnictwo Naukowo-Techniczne. Warszawa.
- Smoluk A. (1998). *O pomiarze, czyli próba próby* [On measurement or how to test a sample]. "Ekonomia Matematyczna" No. 2. Wydawnictwo Akademii Ekonomicznej. Wrocław.
- Wilkowski A. (1998). *Partycje a systemy monetarne* [Partitions and monetary systems]. "Ekonomia Matematyczna" No. 2. Wydawnictwo Akademii Ekonomicznej. Wrocław.
- Wilkowski A. (2000). *O partycjach* [On partitions]. "Ekonomia Matematyczna" No. 4. Wydawnictwo Akademii Ekonomicznej. Wrocław.
- Wilkowski A. (2007). *Uwagi o prawdopodobieństwie częstościowym i przestrzeni probabilistycznej* [Remarks on frequency probability and probabilistic space]. "Didactics of Mathematics" No. 4 (8). Wydawnictwo Akademii Ekonomicznej. Wrocław.

Żabiński Z. (1989). *Rozwój systemów pieniężnych w Europie Zachodniej i Północnej* [Development of monetary systems in North-West Europe]. Ossolineum. Wrocław.